Preparing and purifying an organic liquid

AQA 7404/7405 · AS content

Answer on your own paper. Show your working where needed.

1. Choose an apparatus arrangement

A reaction must be heated for 30 minutes and the starting liquid is volatile. Explain how an upright condenser lets the student keep the mixture hot for that time without losing much starting liquid.

2. Check a thermometer reading

A student sets up simple distillation but puts the thermometer bulb into the boiling liquid. Where should the bulb sit? Explain why the new position gives a more useful reading for identifying the fraction.

3. Check the water connections

A water-cooled condenser is fitted for distillation. Which connector takes in cold water and which lets it out? Explain the direction.

4. Decide which layer to collect

A mixture forms two immiscible layers. The organic layer has density 0.81 g cm^{-3}, and the aqueous layer has density 1.04 g cm^{-3}. The product is in the organic layer. Explain which layer to keep and how to collect it without mixing the layers.

5. Explain the use of a solid

After separating the bulk aqueous layer, a student adds anhydrous magnesium sulfate to the organic layer and later removes the solid. Explain why the salt is used.

6. Interpret the collected fraction

At the same pressure, a pure reference liquid boils at 81 °C. A student collects a fraction from 72 °C to 86 °C. What does this broad range suggest, and what follow-up would improve the evidence?

7. Calculate a preparation yield

A student heats 10.0 g cyclohexanol ($C_6H_{12}O$) with an acid catalyst and isolates 5.74 g cyclohexene (C_6H_{10}). Water is the other product. Write a balanced equation and calculate the percentage yield. Use A_r values C = 12.0, H = 1.0 and O = 16.0.

8. Check a recorded yield

The reaction can give a maximum of 4.60 g pure organic product. A student records collecting 5.08 g of liquid. Calculate the apparent percentage yield. Give one plausible reason for the result and a suitable check.

9. Extra challenge · Plan the collection and judge the yield

A student oxidises 9.00 g of propan-1-ol to propanal, then collects 5.22 g of the aldehyde. The M_r values of propan-1-ol and propanal are 60.0 and 58.0.

Describe how to collect propanal during the reaction and avoid producing mainly propanoic acid. Calculate the percentage yield and explain why it may differ from the percentage of alcohol that reacted.

Scan to try fresh questions online

Try a core or harder question. Use clues and worked answers.

chemistryhelp.co.uk/worksheets/aqa-organic-preparation-and-purification