Polyesters and polyamides

OCR A H432 · A level

Show your working where needed.

Name _____ Date _____

1. Read two polymer chains

Chain A contains …-CH_2CH_2-O-C(=O)-CH_2-C(=O)-O-CH_2CH_2-…

Chain B contains …-CH_2CH_2-NH-C(=O)-CH_2-C(=O)-NH-CH_2CH_2-…

Copy the linking group in each chain. Name each link and classify its polymer.

2. From two monomers to one repeat

Write one repeat unit of the polyester formed from $HOCH_2CH_2OH$ and $HOOCCH_2CH_2COOH$. Show the continuation bond at each end.

3. Find a diamine and diacid

Use the polyamide repeat -[NH-CH_2-CH_2-NH-C(=O)-CH_2-C(=O)]-. Give condensed structures for the diamine and dicarboxylic acid that could make it.

Scan to try fresh questions online

Try a core or harder question. Use clues and worked answers.

chemistryhelp.co.uk/worksheets/ocr-a-polyesters-and-polyamides

Polyesters and polyamides

OCR A H432 · A level

4. Two polymerisation reactions

Propene gives polymer P. A mixture of a diol and a dicarboxylic acid gives polymer Q. Explain what happens to the reacting bonds or groups, name each type of polymerisation and state any other product.

5. Calculate a repeat-unit mass

$HO(CH_2)_3OH$ has M_r 76 and $HOOC(CH_2)_2COOH$ has M_r 118. Calculate M_r for one long-chain polyester repeat containing one fragment of each monomer. Use $M_r(H_2O) = 18$ and ignore end groups.

6. Use alkaline hydrolysis

Give both organic products from complete hydrolysis of -[O-CH_2-CH_2-O-C(=O)-CH_2-CH_2-C(=O)]- with hot excess aqueous NaOH. Show the charge or counter-ions of any salt.

Scan to try fresh questions online

Try a core or harder question. Use clues and worked answers.

chemistryhelp.co.uk/worksheets/ocr-a-polyesters-and-polyamides

Polyesters and polyamides

OCR A H432 · A level

7. Use acidic hydrolysis

Draw the organic products of completely hydrolysing -[NH-CH_2-CH_2-NH-C(=O)-CH_2-C(=O)]- with excess hot dilute HCl. Include any charges in the final mixture.

8. Make a polymer from two different precursors

Use a diacyl chloride and a diamine to make a polyamide with repeat -[NH-$(CH_2)_4$-NH-C(=O)-CH_2CH_2-C(=O)]-. Give both monomers and the small product.

9. Extra challenge · Identify monomers from hydrolysis evidence

A polyamide made from a diamine and a dicarboxylic acid is completely hydrolysed in hot dilute HCl. One product has two carboxylic acid groups separated by three CH_2 groups. The other contains three carbons in a row and two terminal ammonium groups.

Give both products with charges and draw the repeat unit. Predict the form of each product if excess aqueous NaOH had been used instead.

Scan to try fresh questions online

Try a core or harder question. Use clues and worked answers.

chemistryhelp.co.uk/worksheets/ocr-a-polyesters-and-polyamides